



# Digestive Disease Associates

*Specialists in diagnosing and treating gastrointestinal, biliary and liver disorders*

34

10710 Charter Dr., Ste 110  
Columbia, MD. 21044  
(410) 992-9797

700 Geipe Road Suite 230  
Baltimore, MD. 21228  
(410) 247-7500

www.ddamd.com

**Patient:**

**Today's Date:**

**Physician:** Thomas W Abernathy MD

Page 1

## INFORMATION REGARDING YOUR SCHEDULED PROCEDURE MOVIPREP SPLIT DOSE WITH CITRATE OF MAGNESIUM

Date of Procedure: \_ Procedure Time: \_\_\_\_ am/pm Arrival Time: \_\_\_\_ am/pm

Facility:

- ( ) Gastrointestinal Diagnostic Center (410) 242-3636  
700 Geipe Road Suite 220 Baltimore, MD 21228
- ( ) St. Agnes Hospital – GI Services Dept. (410) 368-2200  
900 Caton Ave Baltimore, MD 21229
- ( ) Howard County Gastrointestinal Diagnostic Center (410) 772-7345  
10710 Charter Drive – Suite 120 Columbia, MD 21044
- ( ) The Center for Ambulatory Surgery at Howard County (410) 740-7795  
General Hospital  
5755 Cedar Lane Columbia, MD 21044

General Information:

- IV sedation or anesthesia is usually given during your procedure. **For your safety and due to regulatory requirements, you must arrange for a responsible adult to drive you home.**
- You will not be able to drive, operate machinery, make important decisions or return to work for the rest of the day.
- You may resume normal activities the next day unless the doctor states otherwise.
- If you have any questions concerning your procedure or need to cancel, please call the respective office.
- Wear comfortable, loose fitting clothing that is easy to step into. Wear flat shoes or tennis shoes.
- Do not wear jewelry or bring valuables with you to the Center or hospital.
- **Do not eat or drink anything the day of your procedure or your case will be cancelled.**

## **Patient Information - For ALL Gastroenterology Procedures**

### **IF YOU HAVE DIABETES**

- Inform your primary care physician that you will be on clear liquids the day prior to your procedure
- Check your blood sugar frequently while taking the prep and on the morning of your procedure

### **On the day before your procedure**

- Do not take any diabetes pills such as Actos, Glipizide, Glyburide or Metformin
- Reduce your morning and evening doses of long acting insulin by one-half (1/2)

### **On the day of the procedure**

- Do not take any diabetes pills
- Do not take any insulin unless you are using a sliding scale
- It is ok to eat hard candy or drink a clear soft drink if your blood sugars are too low

### **IF YOU ARE ON ASPIRIN, COUMADIN, WARFARIN, PLAVIX, HEPARIN, LOVENOX, FISH OIL OR ANY OTHER BLOOD THINNERS**

- Your gastroenterologist may have advised you to stop taking blood pressure thinners prior to your procedure. **You will need to stop taking these medications 5 - 7 days prior to your procedure**, based on the recommendations of your gastroenterologist.
- Check with your cardiologist or primary care physician (or the anti-coagulation clinic) if these medications have been prescribed for you due to a heart condition or because you are at a high risk for heart attack or stroke

### **INFORMATION IF YOU TAKE ASPIRIN**

If you take aspirin, it is necessary for you to stop taking aspirin and any anti-inflammatory arthritis medications (such as Advil, Motrin, Aleve and Naprosyn). You should stop seven (7) days prior to the procedure, unless you are at high risk for heart attack or stroke. Check with your doctor about discontinuing your aspirin and plavix. Tylenol is OK to take.

### **WHAT IS A CLEAR LIQUID DIET?**

As a rule – if you can see through it, you can drink it – water is great. Gatorade is an alternative to water though you should avoid drinking red or orange Gatorade. Note that if you are a diabetic or have kidney disease, water is recommended.

### **ITEMS THAT YOU CAN EAT OR DRINK**

- Juices - white grape juice, apple juice, or white cranberry juice.
- Kool-Aide, Powerade or Crystal Light (**no red, blue, orange, or purple**)
- Popsicles (**no red, blue, orange, or purple**)
- Tea or coffee (no milk, cream or non-dairy products) Sugar is ok
- Soda (**no red, blue, orange, or purple**) 7-Up, Sprite, or Ginger Ale
- Broth or bouillon: Beef, Chicken or Vegetable flavored
- Hard candies
- Jell-O (**no red, blue, orange, or purple**)

If you have Diabetes or on Coumadin, Warfarin, Plavix, Heparin, Fish Oil, Lovenox, or any other anticoagulants, please follow the below instructions:

**Preparation Instructions for Colonoscopy – Moviprep Split Dose**

(Please only follow these instructions and pay attention to the following dates)

<p><b>7 Days</b> before your colonoscopy</p>	<ul style="list-style-type: none"> <li>• Read all prep instructions</li> <li>• Contact prescribing physician for instructions on dosage of blood thinners.</li> <li>• Stop herbals, vitamins and oral iron supplements.</li> <li>• Avoid any salads or high fiber foods including nuts, seeds and popcorn.</li> <li>• Stop any anti-inflammatory medications (Motrin, Advil, and Ibuprofen). Tylenol is OK to use.</li> <li>• Arrange for a driver to take you home after the procedure.</li> </ul>
<p><b>2 Days</b> before your colonoscopy</p>	<ul style="list-style-type: none"> <li>• List all current medications.</li> <li>• Obtain bowel prep prescription and a 10 oz bottle of Citrate of Magnesia from your pharmacy. <b>The MoviPrep carton</b> contains 4 pouches and a disposable container for mixing</li> <li>• Take the 10 oz bottle of Citrate of Magnesia after your last meal that night.</li> <li>• Obtain two 32 ounce bottles of Gatorade, Sprite or Ginger Ale. No red or purple fluids please.</li> <li>• If you are a diabetic, follow the instructions on the previous page and use water instead of Gatorade.</li> </ul> <p><b>YOU ARE TO HAVE NO SOLID FOODS/MEALS AFTER DINNER 2 DAYS BEFORE PROCEDURE</b></p>
<p><b>The day</b> <b>before</b> your colonoscopy</p>	<ul style="list-style-type: none"> <li>• No solid food and no alcohol</li> <li>• Drink plenty of clear liquids all day</li> <li>• Drink an extra 8 ounces of clear liquid every hour while awake. <b>Gatorade is preferred for patients without kidney disease or diabetes.</b></li> <li>• Empty 1 Pouch A and 1 Pouch B into the disposable container.</li> <li>• Add drinking water to the top line on the container. Mix to dissolve and then refrigerate.</li> </ul> <p><b>5 p.m.</b> Every 15 minutes, drink the solution down to the next mark until the full liter is consumed (note that the MoviPrep container is divided by 4 marks).</p> <ul style="list-style-type: none"> <li>• Continue to drink at least 16 - 32 more ounces of clear liquids until you go to bed.</li> <li>• Do not have anything by mouth after midnight.</li> <li>• Apply a petroleum based product or diaper rash ointment to the rectal area if you experience discomfort from frequent stools.</li> <li>• Confirm that you have a driver.</li> </ul>

<p><b>The day of your colonoscopy</b></p>	<ul style="list-style-type: none"> <li>• No solid food and no alcohol</li> <li>• You should take your morning medications (i.e. blood pressure meds) with a small sip of water.</li> </ul> <p><b>5 - 7 hours</b> before the time your procedure is scheduled, empty 1 Pouch A and 1 Pouch B into the disposable container. Add drinking water to the top line on the container and mix to dissolve.</p> <p>Every 15 minutes, drink the solution down to the next mark until the full liter is consumed.</p> <ul style="list-style-type: none"> <li>• If your procedure is scheduled in the early morning, you'll need to get up at night to take this second regimen.</li> <li>• <b>The correct timing of this dose is essential to an effective preparation.</b></li> </ul> <p>After the procedure you may eat your usual diet unless otherwise instructed. Drink 8 ounces of liquid at least 6 times after the procedure and before retiring for the night.</p>
---	---

**NOTE:** You must take ALL of the PREP solution, as directed to clean your bowel adequately. If you are passing clear liquid you must continue to take the entire dose of the prep the day before your exam. If you think the prep is not working, call your physician.

This information is not intended to diagnose health problems or to take the place of medical advice or care you receive from your physician or other health care professional. If you have persistent health problems, or if you have additional questions, please consult your doctor. If you have questions or need more information about your medication, please speak to your pharmacist.

Digestive Disease Associates does not endorse any brand names; any similar products may be used.