DIGESTIVE HEALTH CENTER

Flexible Sigmoidoscopy – Fleets Enema Prep

PLEASE READ ALL INSTRUCTIONS CAREFULLY WHEN PROCEDURE IS SCHEDULED

Flexible Sigmoidoscopy – This procedure involves guiding a flexible, lighted tube-like instrument into the rectum and lower colon.

THE DAY OF PROCEDURE
You may have clear liquids until 2 hours before your exam.
CLEAR LIQUIDS: Must be something you can see through. No red or purple! Examples: Gatorade or Powerade, clear broth or bouillon, coffee, tea (no added milk or creamer), strained fruit juices without pulp, Jell-O, popsicles, Kool-Aid or Crystal Light
*** If you are planning on having sedation – you must stop clear liquids 3 hours before your exam.
 If you are Insulin dependent, do not take your morning dose of insulin. You must check your blood sugar before leaving home. If your level is above 300, please take your Regular insulin according to your sliding scale. Please bring your insulin with you to procedure center.
All patients may take morning medications with sips of water.
At: 2 hours prior to your scheduled procedure time: Take two (2) Fleets laxative type enemas. Follow the instructions on the package.
REMEMBER – if you are going to be sedated: Drinking liquids within these 3 hours will cause us to cancel or postpone your procedure. **Also no gum, hard candy, mints and tobacco products in these 3 hours. If you are unsure of your colon cleansing, please call the Procedure Center