ACCT: DOB: AGE:

DR: DOS:

PRE-OP PHYSICIAN'S ORDERS

Anesthesia Orders:	
□ Admit patient to pre-operative area	
□ IV: □ Lactated Ringers □ 0.9 % NaCl □ D5LR □ D51/3 @ KVO or	ml/hr
□ Urine Pregnancy Test - all female patients of menstruating age. (Exclude those who have undergone a Hysterectomy	
or Tubal Ligation/Ablation, or menopausal females with absense of menses 1 year or great	start
□ Fasting Blood Glucose for all NIDDM and IDDM patients - Urine Ketones for FBS over 200	
□ Versed Elixirmlmg as pre-medication for pediatric patients	<u> </u>
Anesthesiologist Signature:	
Additional Anesthesia Orders:	
Surgeon Orders:	
□ Shave Operative Site if warranted	
Nerve Block for Post-Operative Pain Control	
□ Anterior Scalene Block of Brachial Plexus □ Left	□ Right
□ Analgesic Nerve Block of Femoral Nerve □ Left	□ Right
□ Caudal Block	La I Ny II
Antibiotics to be given by Anesthesia within 30 minutes of incision time.	
□ Ancef 1 g IV push □ Ancef 2 g IV push	
□ Cleocin 600 mg IV push	
□ Cleocin 900 mg IV push	
□Other	
Surgeon Signature:	7.474.4